




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 409/14, A61K 31/445, A61P 25/24, 25/22</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/40579</b>  <b>(43) International Publication Date:</b> 13 July 2000 (13.07.00)
<b>(21) International Application Number:</b> PCT/US00/00348  <b>(22) International Filing Date:</b> 6 January 2000 (06.01.00)  <b>(30) Priority Data:</b> 09/226,552                      7 January 1999 (07.01.99)                      US  <b>(71) Applicant:</b> AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US).  <b>(72) Inventors:</b> MEWSHAW, Richard, Eric; Apartment B-509, 251 West DeKaib Pike, King of Prussia, PA 19406 (US). BAUDY, Reinhardt, Bernhard; 5281 Harrington Court, Doylestown, PA 18901 (US).  <b>(74) Agents:</b> NAGY, Michael, R.; American Home Products Corporation, Patent Law Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US) et al.		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> NEW 1,4-DISUBSTITUTED CYCLOHEXANE DERIVATIVES FOR THE TREATMENT OF DEPRESSION  <b>(57) Abstract</b>  Compounds useful for the treatment of disorders of the serotonin-affected neurological systems are provided which have formula (I) wherein: X is carbon or nitrogen; the dotted line represents an optional bond which is absent when X = N; R <sub>1</sub> and R <sub>2</sub> are each, independently, hydrogen, halogen, CF <sub>3</sub> , alkyl, alkoxy, or MeSO <sub>2</sub> ; R <sub>3</sub> is hydrogen, halogen, or alkyl; R <sub>4</sub> is hydrogen, alkyl, alkylaryl, or aryl; and R <sub>5</sub> is hydrogen, halogen, CF <sub>3</sub> , CN, carbamide, or alkoxy; or a pharmaceutically acceptable salt thereof.		



(I)

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

## NEW 1,4-DISUBSTITUTED CYCLOHEXANE DERIVATIVES FOR THE TREATMENT OF DEPRESSION

This invention relates to compounds useful for the treatment of diseases  
5 affected by disorders of the serotonin-affected neurological systems, such as  
depression and anxiety. More specifically, the present invention is directed to 1,4-  
disubstituted cyclohexane derivatives for the treatment of such disorders, processes  
for preparing them and pharmaceutical compositions containing them.

### 10 BACKGROUND OF INVENTION


Pharmaceuticals which enhance the neurotransmission of serotonin (5-HT) are  
useful for the treatment of many psychiatric disorders, including depression and  
anxiety. The first generation of non-selective serotonin-affecting drugs operated  
through a variety of physiological means which caused them to possess numerous  
15 undesired side-effects. The more recently prescribed drugs, the selective serotonin  
reuptake inhibitors (SSRIs), act predominately by inhibiting 5-HT, which is released  
at the synapses, from being actively removed from the synaptic cleft via a presynaptic  
serotonin transport carrier. Since SSRIs require several weeks before they exert their  
full therapeutic effect, this 5-HT blockade mechanism cannot fully account for their  
20 therapeutic activity. It is speculated that this two week induction which occurs before  
a full antidepressant effect is observed, is due to the involvement of the 5-HT<sub>1A</sub>  
autoreceptors which suppress the firing activity of 5-HT neurons, causing a  
dampening of the therapeutic effect. Studies suggest that after several weeks of SSRI  
administration, a desensitization of the 5-HT autoreceptors occurs allowing a full  
25 antidepressant effect in most patients. (See, e.g., Le Poul et al., Arch. Pharmacol.,  
352:141 (1995)). Hence, it is believed that overriding this negative feedback by  
using 5HT<sub>1A</sub> antagonists would potentially increase and accelerate the clinical  
antidepressant response. Recent studies by Artigas et al., Trends Neurosci., 19:378-  
383 (1996), suggest a combination of 5-HT<sub>1A</sub> activity and inhibition of 5-HT uptake  
30 within a single molecular entity can achieve a more robust and fast-acting  
antidepressant effect.

- 2 -

The present invention relates to a new class of molecules which have the ability to act at the 5-HT<sub>1A</sub> autoreceptors and concomitantly with the 5-HT transporter. Such compounds are therefore potentially useful for the treatment of depression as well as other serotonin disorders.

5

U.S. Patent No. 5,468,767 reports a series of substituted indoles of the following formula for the treatment of disorders associated with dysfunction in serotonergic neurotransmission, including depression



wherein R<sub>1</sub> is hydrogen or C<sub>1-4</sub> alkyl;

R<sub>2</sub> is C<sub>1-4</sub> alkyl or (CH<sub>2</sub>)<sub>p</sub>Ar;

m is zero or 1;

n is an integer from 1 to 3; and

15 p is zero or an integer from 1 to 4.

U.S. Patent No. 5,622,951 discloses a series of piperazine derivatives of the following formula for the treatment of CNS disorders, including depression



wherein R is hydrogen or one or two lower alkyl groups;

R<sub>1</sub> and R<sub>2</sub> are each the same or different mono- or bicyclic aryl or heteroaryl radicals;

R<sub>3</sub> is hydrogen, one or two of the same or different lower alkyl groups or a

25 spirocycloalkyl group; and


n is 1 or 2 and m is 1 to 3 and the total of n + m is 2, 3 or 4.

- 3 -

PCT Publication No. WO 93/10092 discloses a series substituted 1,3-cycloalkenes and cycloalkanes useful in the treatment of dopaminergic disorders.

## 5 SUMMARY OF INVENTION

The compounds of this invention are arylpiperazinyl-cyclohexyl indole derivatives represented by Formula I:



X is carbon or nitrogen;

10 the dotted line represents an optional bond which is absent if X is N;

R<sub>1</sub> and R<sub>2</sub> are each, independently, hydrogen, halogen, CF<sub>3</sub>, alkyl, alkoxy, or MeSO<sub>2</sub>;

R<sub>3</sub> is hydrogen, halogen, or alkyl;

R<sub>4</sub> is hydrogen, alkyl, alkylaryl, or aryl; and

R<sub>5</sub> is hydrogen, halogen, CF<sub>3</sub>, CN, carbamide, or alkoxy; or a

15 pharmaceutically acceptable salt thereof.

## DETAILED DESCRIPTION OF THE INVENTION

Preferably, the compounds of the present invention are those of Formula I, wherein:

20 X is carbon; and/or

R<sub>1</sub> and R<sub>2</sub> are each, independently, hydrogen or alkoxy; and/or

R<sub>3</sub> is hydrogen; and /or

R<sub>4</sub> is hydrogen; and/or

R<sub>5</sub> is halogen; or a

25 pharmaceutically acceptable salt thereof.

- 4 -

More preferably, the compounds of the present invention are selected from :  
 5-Fluoro-3-((1,4-cis)-4-[4-(3-methoxy-thiophen-2-yl)-3,6-dihydro-2H-pyridin-1-yl]-cyclohexyl)-1H-indole; and


5-Fluoro-3-((1,4-trans)-4-[4-(3-methoxy-thiophen-2-yl)-3,6-dihydro-2H-pyridin-1-yl]cyclohexyl)-1H-indole.

As used herein, the terms "alkyl" and "alkoxy" as a group or part of a group e.g. arylalkyl are meant to include both straight and branched carbon chains containing 1-6 carbon atoms. The term "aryl" is meant to include aromatic radicals of 6-12 carbon atoms. The term "halogen" is meant to include fluorine, chlorine, bromine and iodine.

The compounds of this Formula I also may be used in the form of a pharmaceutically acceptable acid addition salt having the utility of the free base. Such salts, prepared by methods well known to the art, are formed with both inorganic or organic acids, for example: fumaric, maleic, benzoic, ascorbic, pamoic, succinic, bismethylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, oxalic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzene-sulfonic, hydrochloric, hydrobromic, sulfuric, cyclohexylsulfamic, phosphoric and nitric acids.

The compounds of the present invention may be prepared by any suitable method which will be recognized by those skilled in the art.

Accordingly this invention provides a process for preparing compounds of formula I:



(I)

- 5 -

wherein:

X is carbon or nitrogen;

the dotted line represents an optional bond which is absent if X is N;

R<sub>1</sub> and R<sub>2</sub> are each, independently, hydrogen, halogen, CF<sub>3</sub>, alkyl, alkoxy, or MeSO<sub>2</sub>;

5 R<sub>3</sub> is hydrogen, halogen, or alkyl;


R<sub>4</sub> is hydrogen, alkyl, arylalkyl, or aryl; and

R<sub>5</sub> is hydrogen, halogen, CF<sub>3</sub>, CN, carbamide, or alkoxy; or  
pharmaceutically acceptable salts thereof,

which comprises one of the following:

10

a) dehydrating a compound of formula II



(II)


wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are as defined above, to give a compound of formula I  
where X is carbon and the optional bond is present;

15

or


- 6 -

b) reacting a compound of formula



5

with a compound of formula (IV):



in which formulae X, the dotted line,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are as defined above, to give a compound of formula I where X is carbon and the optional bond is present;

10

or

c) acidifying a basic compound of formula I with a pharmaceutically acceptable acid to give a pharmaceutically acceptable salt;

15

or

d) separating a mixture of cis and trans isomers of a compound of formula (I) to isolate one isomer substantially free from the other isomer.

20


With regard to process a) the dehydration may take place by using an acid, e.g. acetic acid and optionally heating, e.g. at about 70 °C.



- 7 -

Process b) may be carried out by reductive alkylation, e.g. using reducing agent such as sodium triacetoxyborohydride in a suitable solvent e.g. acetic acid.

- A compound of formula (III) may be prepared by reaction of an aminothiophene with  
 5 nitrogen mustard (R = H) or an N-protected derivative e.g. R = benzyl:



followed by deprotection. The reaction is conveniently carried out using a high boiling point solvent, e.g. xylene, toluene or butanol, optionally with base, e.g.  $K_2CO_3$ .

10

The compounds of formula I may be isolated in the form of a salt of a pharmaceutically acceptable acid, e.g. an organic or inorganic acid by treatment with an acid such as described above.

- 15 Geometric (cis and trans) isomers are possible and such isomers can be separated by standard techniques eg chromatography.


The starting materials/reactants used in the processes above are known or can be made by methods known in the art from readily available materials by processes

- 20 known or readily apparent to those skilled in the art.

However, the present compounds may be advantageously prepared according to Scheme 1 set forth below.

25

- 8 -

**Scheme 1**

Specific exemplification of the production of representative compounds of this invention is given in the following procedures.

5

**INTERMEDIATE 1****4-(5-Fluoro-1H-3-indolyl)-cyclohex-3-en-ethylene ketal**

5-Fluoroindole (4.96 g, 0.036 mol), 1,4-cyclohexanedione monoethylene ketal (7.17 g, 0.046 mol) and potassium hydroxide (6 g, 0.043 mol) were heated to reflux in 70 ml of methanol for 6 hours. The reaction was cooled and the product was isolated by filtration and washed with water to give 8.59 g (86%) of product as a white solid: mp 153-155°C.

15

**INTERMEDIATE 2****4-(5-Fluoro-1H-3-indolyl)-cyclohexanone ethylene ketal**

A mixture of 4-(5-fluoro-1H-3-indolyl)-cyclohex-3-en-ethylene ketal (8.5 g) and 10% palladium on carbon (2.72 g) in ethanol (200 ml) was hydrogenated for 5 hours. The catalyst was filtered off and the solvent removed under vacuum.

Chromatography (methanol-methylene chloride) afforded 7.55 g (82 %) of product as a white solid; mp 183-185°C.

### INTERMEDIATE 3

5 **4-(5-Fluoro-1H-3-indolyl)-cyclohexanone**

A solution of 4-(5-fluoro-1H-3-indolyl)-cyclohexanone ethylene ketal (2.8 g, 10 mmol) in 2 L (1:1) tetrahydrofuran-hydrochloric acid (1N) was allowed to stir at room temperature for 16 hours. The solvent was evaporated under vacuum. The crude product was dissolved in ethyl acetate, and washed with 1N sodium hydroxide (3 x 150 ml). The organic layer was dried over anhydrous sodium sulfate and filtered. Chromatography (40% ethyl acetate-hexanes) afforded 2.1 g (91%) of product as yellow solid; mp 112-114°C.

## INTERMEDIATE 4

15 **5-Fluoro-3-((1,4-cis)-4-hydroxy-4-[4-(3-methoxy-thiophen-2-yl)-3,6-dihydro-2H-pyridin-1-yl]-cyclohexyl)-1H-indole**

A solution of 4-(5-fluoro-1H-3-indolyl)-cyclohexanone (1.32 g, 5.7 mmol), 4-hydroxy-4-(3-methoxy-thiophen-2-yl)-piperidine, produced according to the procedures set forth in U.S. Patent No. 5,525,600 (0.5g, 2.5 mmol), sodium triacetoxymethylborohydride (1.82 g, 8.6 mmol) and acetic acid (0.65 ml, 11 mmol) in 1,2-dichloroethane (20 ml) was allowed to stir at room temperature overnight. The reaction was quenched with 1N sodium hydroxide (10 ml), extracted with methylene chloride (3 x 60 ml) and washed with brine (3 x 60 ml). The organic layer was dried over anhydrous sodium sulfate and filtered. Chromatography (10% methanol-ethyl acetate) afforded 0.52 g (22%) of product as a white foam; MS EI *m/e* 428 (M<sup>+</sup>).

## INTERMEDIATE 5

**5-Fluoro-3-((1,4-trans)- 4-hydroxy 4-[4-(3-methoxy-thiophen-2-yl)-3,6-dihydro-2H-pyridin-1-yl]-cyclohexyl)-1H-indole**

30 The trans compound was isolated at same time as the cis isomer in 2.5% yield (0.06 g) as a clear oil; MS EI *m/e* 428 (*M*<sup>+</sup>).

- 10 -

**EXAMPLE 1****5-Fluoro-3-((1,4-cis)-4-[4-(3-methoxy-thiophen-2-yl)- 3,6-dihydro-2H-pyridin-1-yl]-cyclohexyl)-1H-indole**

5           A solution of (0.42 g) of Intermediate 4 in 20 ml acetic acid was heated at 70°C for 0.5 hours. The reaction mixture was poured into 100 ml 2.5 N sodium hydroxide, and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and filtered. Chromatography (10% ethyl acetate-hexanes) afforded 0.32 g of product as a yellow oil, MS EI *m/e* 410 (*M*<sup>+</sup>).

10          The HCl salt was prepared in ethyl acetate: mp 64-167°C.

Elemental analysis for C<sub>24</sub>H<sub>27</sub>FOSN<sub>2</sub>•HCl•1.25H<sub>2</sub>O•0.07C<sub>4</sub>H<sub>8</sub>O<sub>2</sub>

Calc'd:       C, 61.39; H, 6.55; N, 5.97

Found:        C, 61.50; H, 6.19; N, 6.02

15

**EXAMPLE 2****5-Fluoro-3-((1,4-trans)-4-[4-(3-methoxy-thiophen-2-yl)- 3,6-dihydro-2H-pyridin-1-yl]-cyclohexyl)-1H-indole**

          This compound was prepared in the manner described above for Example 1 with the exception that Intermediate 5 (0.48 g) in 20 ml acetic acid was heated to provide in 70% (0.32 g) yield of product as a white solid: mp 190.5-191.5°C.

20          The HCl salt was prepared in ethyl acetate: mp 253.5-255.5°C.

Elemental analysis C<sub>24</sub>H<sub>27</sub>FOSN<sub>2</sub>•HCl•

Calc'd:       C, 64.49; H, 6.31; N, 6.27

Found:        C, 64.07; H, 6.22; N, 6.01

25

          The activity of the present compounds is demonstrated by the following standard pharmacological test procedures.

          The PCR cloning of the human 5-HT<sub>1A</sub> receptor subtype from a human genomic library has been described previously by Chanda et al., Mol. Pharmacol., 43:516 (1993). A stable Chinese hamster ovary cell line expressing the human 5-HT<sub>1A</sub>

30

- 11 -

receptor subtype (5-HT<sub>1A</sub>.CHO cells) was employed throughout this study. Cells were maintained in DMEM supplemented with 10% fetal calf serum, non-essential amino acids and penicillin/ streptomycin.

5           Cells were grown to 95-100% confluency as a monolayer before membranes were harvested for binding studies. Cells were gently scraped from the culture plates, transferred to centrifuge tubes, and washed twice by centrifugation (2000 rpm for 10 min., 4°C) in buffer (50 mM Tris; pH 7.5). The resulting pellets were aliquoted and maintained at -80°C. On the day of assay, the cells were thawed on ice, and  
10 resuspended in buffer. Studies were conducted using [<sup>3</sup>H]8-OH-DPAT as the radioligand. The binding assay was performed in 96 well microtiter plates in a final total volume of 250 µL of buffer. Comparison experiments were performed by using 7 concentrations of unlabelled drug and a final ligand concentration of 1.5 nM. Non-specific binding was determined in the presence of 10 µM 5HT. Saturation analysis  
15 was conducted by using [<sup>3</sup>H]8-OH-DPAT at concentrations ranging from 0.3-30 nM. Following a 30 minute incubation at room temperature, the reaction was terminated by the addition of ice cold buffer and rapid filtration using a M-96 Brandel Cell Harvester (Gaithersburg, MD) through a GF/B filter presoaked for 30 minutes in 0.5% polyethyleneimine.

20

A protocol similar to that used by Cheetham et al., Neuropharmacol., 32:737 (1993) was used to determine the affinity of compounds for the serotonin transporter. Briefly, frontal cortical membranes prepared from male Sprague-Dawley rats were incubated with <sup>3</sup>H-paroxetine (0.1 nM) for 60 min at 25°C. All tubes also contained  
25 either vehicle, test compound (one to eight concentrations), or a saturating concentration of fluoxetine (10 µM) to define specific binding. All reactions were terminated by the addition of ice cold Tris buffer followed by rapid filtration using a Tom Tech filtration device to separate bound from free <sup>3</sup>H-paroxetine. Bound radioactivity was quantitated using a Wallac 1205 Beta Plate<sup>®</sup> counter. Nonlinear  
30 regression analysis was used to determine IC<sub>50</sub> values which were converted to Ki

- 12 -

values using the method set forth in Cheng and Prusoff, Biochem. Pharmacol., 22:3099 (1973) ( $K_i = IC_{50}/((\text{Radioligand conc.})/(1 + K_D))$ ).

The [ $^{35}$ S]-GTP $\gamma$ S binding assay was similar to that used by Lazareno and  
5 Birdsall, Br. J. Pharmacol. 109:1120 (1993). Briefly, 5-HT $_{1A}$  cloned receptor  
membrane fragments (as used for 5-HT $_{1A}$  receptor binding assays) were stored at  
-70°C. When needed, membranes were rapidly thawed, centrifuged at 40,000 x g for  
10 minutes and resuspended at 4 °C for 10 minutes in assay buffer (25 mM HEPES, 3  
mM MgCl $_2$ , 100 mM NaCl, 1 mM EDTA, 10 uM GDP, 500 mM DTT, pH 8.0).  
10 These membranes were then incubated for 30 min at 30 °C with [ $^{35}$ S]GTP $\gamma$ S (1 nM)  
in the presence of vehicle, test compound (one to eight concentrations), or excess 8-  
OH-DPAT to define maximum agonist response. All reactions were terminated by  
the addition of ice cold Tris buffer followed by rapid filtration using a Tom Tech®  
filtration device to separate bound from free [ $^{35}$ S]GTP $\gamma$ S. Agonists produced an  
15 increase in the amount of [ $^{35}$ S]GTP $\gamma$ S bound whereas antagonists produced no  
increase in binding. Bound radioactivity was counted and analyzed as above.

The following assays were performed by incubating the cells with DMEM  
containing 25 mM HEPES, 5 mM theophylline and 10  $\mu$ M pargyline for a period of  
20 20 minutes at 37°C. Functional activity was assessed by treating the cells with  
forskolin (1 uM final concentration) followed immediately by test compound (6  
concentrations) for an additional 10 min at 37°C. In separate experiments, 6  
concentrations of antagonist were preincubated for 20 min prior to the addition of 10  
nM 8-OH-DPAT and forskolin. The reaction was terminated by removal of the  
25 media and addition of 0.5 ml ice cold assay buffer. Plates were stored at -20°C prior  
to assessment of cAMP formation by a cAMP SPA assay (Amersham).

The compounds tested correspond to those prepared in Examples 1 and 2  
above. The results of the procedures are set forth in Table 1.

- 13 -

Example No.	5-HT <sub>1A</sub> (K <sub>i</sub> , nM)	ST (K <sub>i</sub> , nM, )
1	48% @ 1000 nM	2.5
2	20.4	18% @ 100 nM

As demonstrated by the results set forth above, the compounds of the present invention are active towards 5HT1A receptors and generally elevate serotonin levels by inhibiting 5-HT transport. Accordingly, the present compounds should be useful in treating disorders related to defects in serotonin concentration.

The compounds of this invention may be administered orally or parenterally, neat or in combination with conventional pharmaceutical carriers. Applicable solid carriers can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents or an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Any of the solid carriers known to those skilled in the art may be used with the compounds of this invention. Particularly suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

Liquid carriers may be used in preparing solutions, suspensions, emulsions, syrups and elixirs of the compounds of this invention. The compounds of this invention can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fat. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and

- 14 -

parenteral administration include water (particularly containing additives as above, e.g., cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) and their derivatives and oils (e.g., fractionated coconut oil and arachis oil). For  
5 parenteral administration, the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used in compositions for parenteral administration.

Liquid pharmaceutical compositions which are sterile solutions or suspensions  
10 can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. Compositions for oral administration may be either liquid or solid composition form.

Preferably, the pharmaceutical compositions containing the compounds of this  
15 invention are in unit dosage form, e.g., tablets or capsules. In such form, the compositions may be sub-divided in unit doses containing appropriate quantities of the present compounds. The unit dosage forms can be packaged compositions, for example, packeted powders, vials, ampoules, prefilled syringes or sachets containing liquids. Alternatively, the unit dosage form can be, for example, a capsule or tablet  
20 itself, or it can be the appropriate number of any such compositions in package form.

The therapeutically effective amount of the compounds of this invention that is administered and the dosage regimen depends on a variety of factors, including the weight, age, sex, and medical condition of the subject, the severity of the disease, the  
25 route and frequency of administration, and the specific compound employed, and thus may vary widely. However, it is believed that the pharmaceutical compositions may contain the compounds of this invention in the range of about 0.1 to about 2000 mg, preferably in the range of about 0.5 to about 500 mg and more preferably between about 1 and about 100 mg. Projected daily dosages of active compound are  
30 about 0.01 to about 100 mg/kg body weight. The daily dose can be conveniently administered two to four times per day.


The present invention may be embodied in other specific forms without departing from the spirit and essential attributes thereof and accordingly, reference  
35 should be made to the appended claims, rather than to the foregoing specification, as indicating the scope of the invention.



- 15 -

## CLAIMS:

1. A compound of the formula:



5 wherein:

X is carbon or nitrogen;

the dotted line represents an optional bond which is absent if X is N;

R<sub>1</sub> and R<sub>2</sub> are each, independently, hydrogen, halogen, CF<sub>3</sub>, alkyl, alkoxy, or MeSO<sub>2</sub>;

R<sub>3</sub> is hydrogen, halogen, or alkyl;

10 R<sub>4</sub> is hydrogen, alkyl, alkylaryl, or aryl; and

R<sub>5</sub> is hydrogen, halogen, CF<sub>3</sub>, CN, carbamide, or alkoxy; or a pharmaceutically acceptable salt thereof.

2. A compound as claimed in claim 1 wherein X is carbon.

15

3. A compound as claimed in claim 1 or claim 2 wherein R<sub>1</sub> and R<sub>2</sub> are each, independently, hydrogen or alkoxy.

4. A compound as in any one of claims 1 to 3 wherein R<sub>3</sub> is hydrogen.

20

5. A compound as in any one of claims 1 to 4 wherein R<sub>4</sub> is hydrogen.

6. A compound as in any one of claims 1 to 5 wherein R<sub>5</sub> is halogen.

- 16 -

7. The compound of claim 1 which is 5-Fluoro-3-((1,4-cis)-4-[4-(3-methoxy-thiophen-2-yl)-3,6-dihydro-2H-pyridin-1-yl]-cyclohexyl)-1H-indole or a pharmaceutically acceptable salt thereof.


5 8. The compound of claim 1 which is 5-Fluoro-3-((1,4-trans)-4-[4-(3-methoxy-thiophen-2-yl)-3,6-dihydro-2H-pyridin-1-yl]cyclohexyl)-1H-indole indole or a pharmaceutically acceptable salt thereof.

9. A pharmaceutical composition comprising a compound of the formula  
10 (I) as claimed in any one of claims 1 to 8 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

10. A method of treating depression in a patient in need thereof comprising administering to said patient an antidepressant effective amount of a  
15 compound of the formula (I) as claimed in any one of claims 1 to 8 or a pharmaceutically acceptable salts thereof.

11. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 8 or pharmaceutically acceptable salt thereof, which comprises one  
20 of the following:

a) dehydrating a compound of formula II




(II)

wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are as defined in claim 1, to give a compound of formula I as defined in claim 1 where X is carbon and the optional bond is present;


25 or

- 17 -

b) reacting a compound of formula



with a compound of formula (IV):



5

in which formulae X, the dotted line,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are as defined in claim 1, to give a compound of formula I as defined in claim 1 where X is carbon and the optional bond is present;

or

- 10 c) acidifying a basic compound of formula I with a pharmaceutically acceptable acid to give a pharmaceutically acceptable salt

or

- d) separating a mixture of cis and trans isomers of a compound of formula (I) to isolate one isomer substantially free from the other isomer.

15

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/00348

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D409/14 A61K31/445 A61P25/24 A61P25/22

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 92 12977 A (WARNER LAMBERT CO) 6 August 1992 (1992-08-06) abstract page 10 -page 11 page 11, line 26 - line 29	1,9,10
Y	WO 94 15928 A (WYETH JOHN & BROTHER LTD ;WARD TERENCE JAMES (GB); ASHWELL MARK AN) 21 July 1994 (1994-07-21) cited in the application abstract; claims 1,8	1,9,10
Y	US 5 468 767 A (CIPOLLINA JOSEPH A ET AL) 21 November 1995 (1995-11-21) cited in the application abstract; claims 7-9	1,9,10
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 May 2000

Date of mailing of the international search report

06/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Paisdor, B

## INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 00/00348

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 93 10092 A (WARNER LAMBERT CO) 27 May 1993 (1993-05-27) page 7, line 24 -page 8, line 32; claims 3,7,9-11	1,9,10
Y	EP 0 465 398 A (LUNDBECK & CO AS H) 8 January 1992 (1992-01-08) abstract; claims 1,4	1,9,10
Y	EP 0 387 603 A (MERCK PATENT GMBH) 19 September 1990 (1990-09-19) abstract; claim 1; example 1	1,9,10
Y	US 5 322 851 A (PERREGAARD JENS K ET AL) 21 June 1994 (1994-06-21) abstract; claims 1,5; examples 4-6	1,9,10
Y	EP 0 733 628 A (LILLY CO ELI) 25 September 1996 (1996-09-25) abstract; claims; example III page 68; example II	1,9,10
Y	EP 0 431 579 A (WARNER LAMBERT CO) 12 June 1991 (1991-06-12) page 5, line 1 - line 40 page 16; example 1	1,9,10
P,A	EP 0 908 458 A (DUPHAR INT RES) 14 April 1999 (1999-04-14) abstract	1,9,10

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 00/ 00348

### Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/00348

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9212977 A	06-08-1992	US 5124332 A	23-06-1992
WO 9415928 A	21-07-1994	AT 155133 T	15-07-1997
		AU 5711594 A	15-08-1994
		DE 69312091 D	14-08-1997
		DK 677046 T	01-09-1997
		EP 0677046 A	18-10-1995
		ES 2105617 T	16-10-1997
		GR 3024430 T	28-11-1997
		IL 108259 A	24-09-1998
		JP 8505381 T	11-06-1996
		US 5622951 A	22-04-1997
US 5468767 A	21-11-1995	US 5607961 A	04-03-1997
WO 9310092 A	27-05-1993	US 5314896 A	24-05-1994
		AU 664872 B	07-12-1995
		AU 3129193 A	15-06-1993
		CZ 9401161 A	16-11-1994
		EP 0545095 A	09-06-1993
		EP 0613465 A	07-09-1994
		FI 942310 A	15-07-1994
		FI 980471 A	02-03-1998
		HU 70520 A	30-10-1995
		JP 7501075 T	02-02-1995
		MX 9206635 A	29-07-1994
		NO 941884 A,B,	19-05-1994
		NZ 245174 A	26-09-1995
		RU 2116297 C	27-07-1998
		SG 46641 A	20-02-1998
		SK 56994 A	08-02-1995
		US 5409931 A	25-04-1995
		ZA 9208955 A	19-05-1994
		MX 9302225 A	29-04-1994
EP 0465398 A	08-01-1992	AT 115576 T	15-12-1994
		AU 644186 B	02-12-1993
		AU 8014491 A	02-01-1992
		CA 2045955 A	03-01-1992
		DE 69105839 D	26-01-1995
		DE 69105839 T	27-04-1995
		DK 465398 T	22-05-1995
		ES 2064973 T	01-02-1995
		FI 913194 A	03-01-1992
		GR 3015286 T	30-06-1995
		HK 52195 A	13-04-1995
		IE 65339 B	18-10-1995
		IL 98495 A	29-06-1995
		JP 4253976 A	09-09-1992
		KR 9510163 B	11-09-1995
		NO 178191 B	30-10-1995
		NZ 238757 A	26-08-1993
		PT 98168 A,B	29-05-1992
		SG 19895 G	18-08-1995
		US 5322851 A	21-06-1994
		US 5457115 A	10-10-1995
		US 5216001 A	01-06-1993
		ZA 9105110 A	27-05-1992

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/00348

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0387603 A	19-09-1990	DE 3907974 A	13-09-1990
		AU 622291 B	02-04-1992
		AU 5116290 A	13-09-1990
		CA 2011834 A	11-09-1990
		HU 56088 A,B	29-07-1991
		JP 2273672 A	08-11-1990
		ZA 9001857 A	28-12-1990
US 5322851 A	21-06-1994	US 5457115 A	10-10-1995
		AT 115576 T	15-12-1994
		AU 644186 B	02-12-1993
		AU 8014491 A	02-01-1992
		CA 2045955 A	03-01-1992
		DE 69105839 D	26-01-1995
		DE 69105839 T	27-04-1995
		DK 465398 T	22-05-1995
		EP 0465398 A	08-01-1992
		ES 2064973 T	01-02-1995
		FI 913194 A	03-01-1992
		GR 3015286 T	30-06-1995
		HK 52195 A	13-04-1995
		IE 65339 B	18-10-1995
		IL 98495 A	29-06-1995
		JP 4253976 A	09-09-1992
		KR 9510163 B	11-09-1995
		NO 178191 B	30-10-1995
		NZ 238757 A	26-08-1993
		PT 98168 A,B	29-05-1992
		SG 19895 G	18-08-1995
		US 5216001 A	01-06-1993
		ZA 9105110 A	27-05-1992
EP 0733628 A	25-09-1996	AU 702322 B	18-02-1999
		AU 5311296 A	08-10-1996
		BR 9601061 A	06-01-1998
		CA 2215322 A	26-09-1996
		CN 1184425 A	10-06-1998
		CZ 9702888 A	18-02-1998
		HU 9800417 A	28-06-1999
		JP 11502816 T	09-03-1999
		NO 974220 A	04-11-1997
		NZ 305166 A	23-12-1998
		PL 322843 A	16-02-1998
		WO 9629075 A	26-09-1996
		US 5708008 A	13-01-1998
		US 5962474 A	05-10-1999
EP 0431579 A	12-06-1991	US 4975445 A	04-12-1990
		AT 109136 T	15-08-1994
		AU 630770 B	05-11-1992
		AU 6770190 A	13-06-1991
		CA 2031570 A	07-06-1991
		DE 69011054 D	01-09-1994
		DE 69011054 T	08-12-1994
		DK 431579 T	31-10-1994
		ES 2057341 T	16-10-1994
		FI 905957 A	07-06-1991
		IE 66117 B	13-12-1995



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/00348

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0431579 A		JP 3251576 A	11-11-1991
		KR 166359 B	15-01-1999
		NO 176097 B	24-10-1994
		NZ 236336 A	28-04-1992
		PH 27326 A	08-06-1993
		PT 96084 A,B	30-09-1991
		ZA 9009774 A	26-08-1992
EP 0908458 A	14-04-1999	AU 8607998 A	15-04-1999
		CA 2247734 A	24-03-1999
		CN 1220265 A	23-06-1999
		CZ 9803030 A	14-04-1999
		JP 11302279 A	02-11-1999
		NO 984380 A	25-03-1999
		PL 328754 A	29-03-1999
		SK 129498 A	13-04-1999